Available online at www.sciencedirect.com

Public Health

journal homepage: www.elsevier.com/puhe



'Loneliness in the city': examining socio-economics, loneliness and poor health in the North East of England



S.J. Macdonald ^{a,*}, J. Nixon ^b, L. Deacon ^c

^a School of Social Sciences, Faculty of Education and Society, University of Sunderland, The Reg Vardy Centre, St Peter's Way, Sunderland, SR6 ODD, UK

^b People Services Directorate – Public Health, Sunderland Civic Centre, Burdon Road, Sunderland City Council, UK ^c Applied Social Studies/Social Work, Faculty of Education and Society, The Reg Vardy Centre, St Peter's Way, Sunderland, SR6 0DD, UK

ARTICLE INFO

Article history: Received 23 March 2018 Received in revised form 24 August 2018 Accepted 5 September 2018

Keywords: Inequalities Socio-economics Annual income Isolation Loneliness Poverty

ABSTRACT

Objectives: The aim of this study was to discover whether lower socio-economic status is associated with increased experiences of loneliness and isolation. The research subsequently determined whether this relationship impacted health inequalities.

Study design: The study used a cross-sectional, self-reported survey collecting information on loneliness, isolation and poor health (n = 680). The survey was administered through Sunderland District Council in 2016–2017, and data were analysed at The University of Sunderland.

Methods: The study used a quantitative approach, and data were analysed using descriptive statistics, engaging in univariate, bivariate and multivariate levels of analysis.

Results: A number of significant findings emerged from the data analysis, linking lower socio-economic status to experiences of loneliness (P = 0.000) and social isolation (P = 0.000). When determining if social isolation and socio-economics had a detrimental impact on a person's health, no statistical association was discovered (P = 0.098). Yet, there was a significant relationship concerning socio-economic status, loneliness and poor health (P = 0.026).

Conclusions: The authors have identified a number of associations within the data with reference to isolation, loneliness and poor health. Therefore, participants from a lower socio-economic group experienced disproportionately high levels of social isolation and emotional loneliness when compared with other socio-economic groups. The data also demonstrate that participants who experienced loneliness, and who were from a lower socio-economic background, were consistently more likely to report poor health than those from other socio-economic backgrounds.

 \odot 2018 The Royal Society for Public Health. Published by Elsevier Ltd. All rights reserved.

* Corresponding author.

https://doi.org/10.1016/j.puhe.2018.09.003

E-mail addresses: stephen.j.macdonald@sunderland.ac.uk (S.J. Macdonald), Jackie.Nixon@sunderland.gov.uk (J. Nixon), lesley. deacon@sunderland.ac.uk (L. Deacon).

^{0033-3506/© 2018} The Royal Society for Public Health. Published by Elsevier Ltd. All rights reserved.

Introduction

The association between health inequalities and the socioeconomic status has been significantly documented over the past three decades.^{1–5} The Marmot review substantiates that employment and economic prosperity directly influence the UK population's health. Inequalities in health arise because of inequalities in the conditions of daily life. The conditions in which people are born, grow, live, work and age are fundamentally affected by inequalities of money, resources and power.^{3,4} The World Health Organisation suggests that our health is determined by a complex interaction between our individual characteristics, lifestyle and the physical, social and economic environment which we occupy.^{6,7} Despite the commitment and implementation of policies from the government in England and Wales, the gap in health inequalities is widening within some geographical regions, particularly in the North of England.^{8,5} After the 2008 banking crisis, austerity measures resulted in increased economic deprivation which had a significant impact within the North East regions of England.9 As Whitehead indicates, communities across the North East of England have seen a rise in health inequalities and premature deaths, when compared with the national average.⁸ As Whitehead⁸ suggests, the City of Sunderland, which is situated within the North East of England, was specifically identified as a geographical area of concern with regard to rising health inequalities. Sunderland has a population of approximately 277,000 and the Indices of Multiple Deprivation¹⁰ indicate that 38% of the city's population live in areas that are among the 20% most disadvantaged across England, with the annual household income significantly lower than the national average. Hence, income is a key determinant of deprivation, and evidence shows that societies with a bigger gap between rich and poor have worse health outcomes overall.3

Although there have been numerous studies linking socioeconomics with health inequalities, very few studies have investigated the intersectional impact that socio-economics and loneliness have on poor health.¹¹ However, defining the concept of loneliness is not without its challenges. Studies in the UK and globally have used different methodologies to collect data on the concept of loneliness. Hence, it can be suggested that the concept of loneliness is a subjective interpretation of a person's day-to-day interactions and is difficult to measure. In previous research, the concepts of loneliness and isolation have been somewhat interconnected and even misinterpreted. Dahlberg and McKee have attempted to clarify the concept of loneliness by referring to the 'emotional' loneliness.²⁰ They suggest that 'emotional loneliness refers to the absence of an attachment figure in one's life and someone to turn to',²⁰ thereby suggesting that 'emotional loneliness' is a subjective interpretation relating to the perceived quality of a person's relationships with others. Furthermore, Tanskanen and Anttila have attempted to clarify this definition further by separating the concepts of loneliness and isolation into different quantitative categories.²¹ They suggest that social isolation and emotional loneliness are distinctly different analytical categories, where the first relates to concrete interactions and the second relates to subjective interpretations

of emotional reactions to a person's environment. Tanskanen and Anttila suggest that 'social isolation is concerned more with environmental impoverishment or restrictions than with the individual's ability to create and maintain social relationships'.²¹ From this perspective, social isolation is underpinned by environmental factors which relate to the breaking down of social networks. Therefore, emotional loneliness can occur due to social isolation because of a lack of contact with family members or friends, but it can also occur even when people do have significant contact with other social groups but where there is a 'lack of desired quality of social engagement'.²¹ From this perspective, an individual can also be socially isolated but not experience feelings of loneliness.

When investigating social isolation and emotional loneliness, a number of studies have illustrated the negative impact this experience has on public health. Previous studies indicate that having no social connections has an increased effect on mortality that exceeds the impact of well-known public health risk factors such as obesity, physical activity and mental wellbeing and has a similar impact to cigarette smoking.^{11–16} As Leigh-Hunt et al. indicate, there is a growing amount of literature linking loneliness and isolation to increasing health inequalities, but very few studies have investigated increased risk factors because of socio-economics.¹¹ Previous studies linking loneliness and isolation to health inequalities have predominantly examined a sample of older and retired participants.^{17,18} Studies on loneliness and isolation that have developed a more inclusive age range sample have predominately reflected on risk factors concerning age and gender, rather than socio-economics per se.^{19–21}

Methods

This article presents findings from a study examining the impact of loneliness and isolation on communities within Sunderland. The study was funded by Sunderland City Council to try to comprehend pathways into loneliness and/or isolation and was conducted to improve and to develop effective services within the district. The research project was conducted over a 1-year period, from September 2016 to September 2017. To collect data on loneliness and isolation, a cross-sectional, mixed-mode method was developed, using a survey that could be completed online or in hard copy.²⁵ It was distributed in a variety of ways to ensure the inclusion of participants from a wide range of social demographics throughout the city, particularly to access socially excluded populations. First, an unrestricted self-selected survey was made available on Sunderland City Council's website.²⁵ This project had been widely publicised throughout the City of Sunderland, and residents could 'opt in' to complete the online survey. However, this approach was supplemented by a team of health visitors, third sector employees and social care professionals, who targeted diverse, and often socially excluded populations across the city.²⁵ Surveys were also made available in local services including, housing, health, criminal justice, libraries, education and immigration. In total, 680 participants took part in the research. Anyone living in the city was invited to take part in the study, whether they identified as experiencing loneliness and/or isolation, and in total, 240 participants self-identified as either experiencing loneliness, isolation or both; this was 35.3% of the sampled population. This allowed the team to make comparisons between the participants who identified as lonely/isolated and participants who did not (N = 440). The data were predominantly coded at the nominal and ordinal level. The team used SPSS to analyse the results using descriptive statistics. Data were analysed by engaging in univariate, bivariate and multivariate analysis.²² The team analysed the data using frequency tables at the univariate level and cross tabulations at the bivariate and multivariate level to produce statistically relevant information.²² At the bivariate and multivariate level, frequency distributions were analysed using a Chi-squared statistic (x^2) or where the expected count reduced below five, a Fisher's Exact Test was used to confirm statistical significance. Data were considered significant if it reduced below the 0.05 statistically significant threshold.²² Only statistically significant data ($P \le 0.05$) are used in the research findings.

Within this article, annual household income was used as the independent variable to discover if socio-economics impacted participants' experiences of emotional loneliness and social isolation.^{20,21} When studying people's experiences of loneliness and social isolation, a number of complexities arose with reference to measurement. As research by Leigh-Hunt et al. illustrates, there are numerous methodological issues when examining previous research on loneliness and isolation.¹¹ In previous research, 'loneliness' has been used as an analytical category that incorporates the aspects of both 'emotional loneliness' and 'social isolation'.^{19,23,24} Drawing on work from Dahlberg and McKee²⁰ and Tanskanen and Anttila,²¹ this study separated the two concepts of emotional loneliness and social isolation. Therefore, the study measured and defined 'emotional loneliness' as a subjective experience, based on strengths of relationships, whereas 'social isolation' refers to a lack of access to social networks within communities. It should be noted that full ethical approval was gained by the research team from Sunderland City Council and the University of Sunderland before the research commenced.

Results

social demographics

At the multivariate stage of analysis, the team discovered that there were no intersectional relationships among age, gender, sexuality and ethnicity with reference to loneliness and isolation. The project attempted to collect data on a wide range of social demographics to examine emotional loneliness and social isolation within Sunderland. Table 1 shows that the sample consisted of a relatively equal age range. The younger category represented participants aged between 17 and 44 years, which consisted of 29% of the sample. The older category of 60 years plus consisted of 34% of the sample. The largest category, at 38%, consisted of the middle-aged group who were aged 45–59 years. Unfortunately, the study did have a gender bias as the majority of participants who completed the study were female at 69%, compared with male at 31%. With reference to sexuality, the vast majority of

Table 1 — Social demographics.										
Social demographic	S	%	Ν							
Sex	Male	30.6%	205							
	Female	69.4%	466							
	Total	100.0%	671							
Age in years	17–29	10.1%	66							
	30-44	18.6%	121							
	45-59	37.5%	244							
	60-74	26.4%	172							
	75+	7.4%	48							
	Total	100.0%	651							
Sexuality	Straight/heterosexual	92.7%	613							
	LGBT	7.3%	48							
	Total	100.0%	661							
Ethnic groups	White	97.8%	655							
	Mixed	0.2%	1							
	Asian	1.0%	7							
	Black	1.0%	7							
	Total	100.0%	670							
Lonely	No	70.9%	482							
	Yes	29.1%	198							
	Total	100.0%	680							
Socially isolated	No	85.0%	578							
	Yes	15.0%	102							
	Total	100.%	680							
Socio-economics:	Below £10,000	26.6%	158							
annual household	£10,000-24,999	34.7%	206							
income	£25,000-49,999	28.2%	167							
	£50,000-100,000	9.1%	54							
	More than £100,000	1.3%	8							
	Total	100.0%	593							
LGBT, lesbian, gay, bis	exual, and transgender.									

the group at 93% identified as heterosexual, compared with lesbian, gay, bisexual, and transgender (LGBT) at 7%. This is consistent with the reported demographics of Sunderland, where 6% of the population comes from a sexual minority background.¹⁰ Furthermore, the vast majority of participants (98%) identified as being from a white ethnic group, with only 2% reporting they were from an ethnic minority background. Again, this is consistent with the ethnic demographics of Sunderland, which has a relatively small ethnic minority population at 4%.¹⁰

Socio-economics, loneliness and isolation

When examining an association among socio-economics (n = 593), experiences of loneliness (n = 198) and social isolation (n = 102; see Table 1), a number of significant findings emerged from the data analysis. This study uncovered a correlation between annual income and social isolation (P = 0.000). In Table 2, the principal group, at 36%, who considered themselves as socially isolated, were people who had an annual household income of less than £10,000. Reports of social isolation declined considerably as household income increased. Only 11% of participants who earned between £10,000–24,999 reported experiencing social isolation. Experiences of social isolation gradually decreased to 4% for people in the £50,000–100,000 group. However, there was a slight increase to 13% for the £100,000+ group. When comparing annual household income of less than £10,000 with that of

Table 2 – Socio-economics and isolation.												
		Isola	ited		Т	'otal	Sig.					
]	No		Yes								
	n	%	n	%	n	%						
Below £10,000	101	63.9%	57	36.1%	158	100.0%	P = 0.000					
£10,000-24,999	183	88.8%	23	11.2%	206	100.0%						
£25,000-49,999	153	91.6%	14	8.4%	167	100.0%						
£50,000-100,000	52	96.3%	2	3.7%	54	100.0%						
More than £100,000	7	87.5%	1	12.5%	8	100.0%						

 $\pm 100,000+$, a reduction can be observed from 36% to 13% from the lowest to the highest socio-economic group.

With reference to the socio-economic status and participants who reported experiencing loneliness, a significant relationship appeared within the data analysis (P = 0.000). As can be seen in Table 3, a staggering 60% of participants whose annual household income was below £10,000 described feelings of emotional loneliness. The experience of loneliness reduced from 60% to 27% for participants whose annual income was from £10,000-24,999. There was a slight variation between the £25,000-49,999 and the £50,000-100,000 group, as loneliness decreased to 13% for the £25,000-49,999 group and subsequently increased to 20% for the £50,000-100,000 group. No participant in the £100,000+ group reported feelings of loneliness. When comparing data for participants with an annual household income of less than £10,000 with those in the $\pm 100,000+$ group, there was a reduction from 60% to 0% between the lowest and highest socio-economic groups.

Risk factors, loneliness and isolation

When exploring risk factors to loneliness and isolation, the team examined if participants described spending most of their time alone. The data presented in Table 4 revealed a significant relationship between socio-economic groups and limited social engagement (P = 0.000). For participants in the below £10,000 group, 68% spent the majority of time on their own. This decreased to 41%, 29% and 17% for the £10,000–24,999 group, £25,000–49,999 group and £50,000–100,000 group, respectively. Interestingly, there was an increase for participants whose household income was £100,000+ as 38% of this group reported spending most of their time on their own. Although there was an increase for this group, when comparing time spent alone for the below £10,000 group with the more than £100,000 group, there is a reduction from 68% to 38% from the lowermost socio-economic groups in this study.

Table 3 – Socio-economics and loneliness.												
		Lone	ely		T	'otal	Sig.					
		No		Yes								
	n	%	n	%	n	%						
Below £10,000	64	40.5%	94	59.5%	158	100.0%	P = 0.000					
£10,000-24,999	151	73.3%	55	26.7%	206	100.0%						
£25,000-49,999	145	86.8%	22	13.2%	167	100.0%						
£50,000-100,000	43	79.6%	11	20.4%	54	100.0%						
More than £100,000	8	100.0%	0	0.0%	8	100.0%						

Table 4 — Time spent alone.												
	Tii	me spe	ent a	llone	Т	'otal	Sig.					
	3	<i>l</i> es	I	No								
	n	%	n	%	n	%						
Below £10,000	102	67.5%	49	32.5%	151	100.0%	P = 0.000					
£10,000-24,999	76	40.6%	111	59.4%	187	100.0%						
£25,000-49,999	42	29.0	103	71.0%	145	100.0%						
£50,000-100,000	9	17.0%	44	83.0%	53	100.0%						
More than £100,000	3	37.5%	5	62.5%	8	100.0%						

Within this study, the team recognised that participants may choose to spend the majority of their time alone. Therefore, the team explored whether participants actively decided to spend their time alone or wanted increased interactions/activities with others. The data in Table 5 revealed that there was a significant correlation between socioeconomic groups and the desire to have increased and reduced contact time with others (P = 0.001). The data revealed that 52% of participants who had an annual income below £10,000 desired to engage in more activities than they currently do outside their households. This decreased to 32% for participants who had an annual income of £10,000-24,999. Interestingly, there was a slight increase to 36% for individuals from the £25,000–49,999 group, but this reduced to 29% for the £50,000-100,000 group. Again, no participants in the £100,000+ group desired any increase in activities and seemed content with their current social networks. The data seem to indicate that for individuals from the lowermost socioeconomic group, there is a desire for increased activities/interactions outside the home, which is not the case for individuals from the uppermost socio-economic group.

Health issues, loneliness and isolation

When comparing the relationship between socio-economics and self-reported poor health, the data revealed that there was a significant correlation between poverty and health in this survey (P = 0.000). Table 6 illustrates that participants who earned less than £10,000 per annum reported the highest level of poor health, at 25%. Furthermore, only 33% of this group described themselves in good health. For participants in the £10,000–24,999 group, only 11% described themselves as in poor health, and 60% of this group described themselves as in good health. For the £25,000–49,999 group, again we could see a reduction in poor health, to 6%. There was also an increase

Table 5 – Desired increase in social interaction.												
		esired i ocial ir		ease in action	Т	'otal	Sig.					
		Yes		No								
	n	%	n	%	n %							
Below £10,000	70	51.5%	66	48.5%	136	100.0%	P = 0.000					
£10,000-24,999	54	32.1%	114	67.9%	168	100.0%						
£25,000-49,999	51	36.4%	89	63.6%	140	100.0%						
£50,000-100,000	14	28.6%	35	71.4%	49	100.0%						
More than £100,000	0	0.0%	8	100.0%	8	100.0%						

Table 6 – Socio-economics and poor health.													
		Socio-e	Т	'otal	Sig.								
	G	ood]	Fair		Bad							
	n	%	n	%	n	%	n	%					
Below £10,000	50	32.5%	66	42.9%	38	24.7%	154	100.0%	P = 0.000				
£10,000-24,999	117	60.0%	56	28.7%	22	11.3%	195	100.0%					
£25,000-49,999	118	76.6%	27	17.5%	9	5.8%	154	100.0%					
£50,000-100,000	43	81.1%	9	17.0%	1	1.9%	53	100.0%					
More than £100,000	6	75.0%	2	25.0%	0	0.0%	8	100.0%					

	Not Lonely										Lonely								
	Socio-economics, poor Total health and loneliness							Socio-economics, poor health and loneliness						Total					
	G	ood	•	Fair	Bad				C	Good		Good Fair		Fair	Bad				
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%			
Below £10,000	30	50.0%	19	31.7%	11	18.3%	60	100.0%	20	21.3%	47	50.0%	27	28.7%	94	100.0%			
£10,000-24,999	96	68.6%	40	28.6%	4	2.9%	140	100.0%	21	38.2%	16	29.1%	18	32.7%	55	100.0%			
£25,000-49,999	107	81.1%	19	14.4%	6	4.5%	132	100.0%	11	50.0%	8	36.4%	3	13.6%	22	100.0%			
£50,000-100,000	38	90.5%	4	9.5%	0	0.0%	42	100.0%	5	45.5%	5	45.5%	1	9.1%	11	100.0%			
More than £100,000	6	75.0%	2	25.0%	0	0.0%	8	100.0%	0	0.0%	0	0.0%	0	0.0%	0	100.0%			
Sig.	P = 0.000 $P = 0.026$																		

in good health, at 77% for this group. Interestingly, only 2% of the \pm 50,000–100,000 group, and none of the \pm 100,000+ group, described themselves as in poor health. Most participants in these groups also described themselves as relatively healthy, as 81% of the \pm 50,000–100,000 group and 75% of the \pm 100,000+ group reported being in good health. Therefore, the data in Table 6 demonstrate that poor health diminishes as household income increases, in this survey.

When comparing the socio-economic status and health inequalities with loneliness and isolation, the data analysis indicated that there was no significant relationship among socio-economics, poor health and social isolation (P = 0.098). However, there was a significant relationship (P = 0.026) among socio-economics, poor health and experiences of loneliness. It should be noted that loneliness is associated with poor health, but experiences of loneliness disproportionately affected the health of people from lower socioeconomic groups. The data in Table 7 demonstrate that, for participants from the below £10,000 group, 29% of participants who had experienced loneliness also reported poor health, compared with 18% of this group who had not experienced loneliness but reported health problems. Interestingly, it was the £10,000-24,999 group where loneliness had the biggest impact, as 33% of participants who had experienced loneliness reported poor health, compared with only 3% of this group who had not experienced loneliness but reported health problems. Poor health then decreased as income increased for people who experienced loneliness. This can be seen in the £25,000-49,999 group who experienced loneliness, as only 14% reported poor health, and 5% of this group who had not experienced loneliness reported health problems. Similarly, only 9% of the £50,000-100,000 group described experiences of loneliness and poor health,

and none of this group who had not experienced loneliness reported having poor health. Finally, none of the $\pm 100,000+$ group reported feelings of either loneliness or experiences of poor health, in this study.

Discussion and Conclusions

The findings seem to illustrate that socio-economics and emotional loneliness affected poor health in this sampled population (P = 0.026). As the findings illustrated, participants with a household annual income of below £10,000 experienced disproportionately high levels of social isolation and emotional loneliness, when compared with other socioeconomic groups (P = 0.000). The group with a household income below £10,000 also spent most of their time on their own, compared with other groups within this survey (P = 0.000). Yet this group was more likely to want increased access to activities outside of their households than the other socio-economic groups within the study. Although the other groups reported spending a significant amount of time on their own, they were less likely to want increased access to more social activities (P = 0.001). This may suggest that individuals who experienced social isolation in the higher socioeconomic groups have had the ability to choose whether they wanted increased or decreased social activities, which is not the case for participants in the below £10,000 group. Therefore, social isolation seems more likely to be a choice for participants as their household income increases, which, to a certain extent, may lead to a lower level of emotional loneliness among these groups.

Finally, when examining if social isolation has a detrimental impact on a person's health, no association was discovered with reference to socio-economics (P = 0.098). However, there was a significant relationship among the socio-economic status, emotional loneliness and poor health (P = 0.026). As the data demonstrated, participants with a household income of less than £24,999 annually were most likely to report poor health compared with the other groups. One might theorise that if participants have no control over their feelings of loneliness, this may have a detrimental impact on their health, whereas individuals who have more choice over their reduced contact with people outside their home have improved health, meaning overall improved health for groups with access to increased economic wealth. Although the article does not present any conclusive findings, it does illustrate a number of noteworthy relationships among socio-economics, loneliness and perceived poor health. Therefore, the findings present data that may be useful as a foundation for future research, examining the complex intersectional relationships among socio-economics, experiences of loneliness and health inequalities.

Strengths and limitations

It should be recognised that no definitive conclusions can be drawn from this article as there were a number of limitations to the survey. First, it should be noted that this study was exploratory in nature and aimed to target a relatively hidden and hard-to-reach community. The survey was conducted partially online, and although this gave the authors the access to a greater number of hard-to-reach participants, it did not allow us to create a sample frame or randomly select a representative population. Second, our socio-economic categories were not evenly distributed within our sample, leading to small numbers in the higher socio-economic groups, which could lead to bias in our findings. Third, information on loneliness and isolation was self-reported, and no objective measurement was used to categorise individuals as either lonely or isolated. Fourth, there was a gender bias within the data collected as 70% of participants were female. Finally, the survey had limited information on participants' socio-economic profiles; hence household income was used as the determinant factor to categorise groups. Although these limitations exist, which may not lead to a precise estimation of the impact that loneliness has on health, the data did present some interesting findings that can be used to design a larger confirmatory study.

Author statements

Ethical approval

Full ethical approval was gained by the research team from Sunderland City Council and the University of Sunderland before the research commenced.

Funding

This research was supported by a small grant from Sunderland City Council. The content of this article is solely the responsibility of the authors and does not represent the official views of Sunderland City Council.

Competing interests

None declared.

REFERENCES

- 1. Marmot MG, Wilkinson RG. Social determinants of health. Oxford: Oxford University Press; 1999.
- Pevalin DJ. Socio-economic inequalities in health and service utilization in the London Borough of Newham. Publ Health 2007;121:596–602.
- **3.** Wilkinson R, Pickett K. The spirit level: why equality is better for everyone. London: Penguin Books; 2009.
- Marmot M. Fair society, healthy lives: the Marmot review executive summary; strategic review of health inequalities in England post-2010. London: Public Health England and UCL Institute of Health Equity; 2010.
- Tweed EJ, Allardice GM, McLoone P, Morrison DS. Socioeconomic inequalities in the incidence of four common cancers: a population-based registry study. *Publ Health* 2018;154:1–10.
- 6. WHO. The role of intersectoral co-operation in national strategies of health for all. Geneva: World Health Organization; 1986.
- Blakely T, Hales S, Woodware A. Assessing the distribution of health risks by socioeconomic position at national and local levels. Geneva: World Health Organization; 2004.
- 8. Whitehead M. Due North, report of the inquiry on health equity for the North. Liverpool: University of Liverpool and Centre for Local Economic Strategies; 2014.
- Clayton J, Donovan C, Merchant J. Distancing and limited resourcefulness: third sector service provision under austerity localism in the north east of England. Urban Stud 2016;53(4):723–40.
- Office for National Statistics. 2014-based subnational population projections for local authorities in England. London: Office for National Statistics; 2016.
- Leigh-Hunt N, Bagguley D, Bash K, Turner V, Turnbull S, Valtorta N, et al. An overview of systematic reviews on the public health consequences of social isolation and loneliness. Publ Health 2017;152:157–71.
- Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. PLoS Med 2010;7(7):e1000316. https://doi.org/10.1371/ journal.pmed.1000316.
- Steptoe A, Shankar A, Demakakos A, Wardle J. Social isolation, loneliness, and all-cause mortality in older men and women. PNAS Direct Submissions 2013;110(15):5797–801.
- Stickley A, Koyanagi A, Leinsalu M, Ferlander S, Sabawoon W, McKee M. Loneliness and health in eastern Europe: findings from moscow, Russia. Publ Health 2015;129:403–10.
- **15.** Watson J, Crawley J, Kane D. Social exclusion, health and hidden homelessness. *Publ Health* 2016;**139**:96–102.
- 16. Elovainio M, Hakulinen C, Pulkki-Raback L, Virtanen M, Josefsson K, Jokela M, et al. Contribution of risk factors to excess mortality in isolated and lonely individuals: an analysis of data from the UK Biobank cohort study. Lancet Publ Health 2017;2.
- Cattan M, White M, Bond J, Learmouth A. Preventing social isolation and loneliness among older people: a systematic review of health promotion intervention. *Aging Soc* 2005;25:41–67.
- Davidson S, Rossall D. Age UK loneliness evidence review. London: Age UK; 2014.
- Victor CR, Bowling A. A longitudinal analysis of loneliness among older people in great Britain. J Psychol 2012;146(3):313–31.

- 20. Dahlberg L, McKee KJ. Correlates of social and emotional loneliness in older people: evidence from an English community study. *Aging Ment Health* 2014;**18**(4):504–14.
- Tanskanen J, Anttila T. A prospective study of social isolation, loneliness, and mortality in Finland. Am J Publ Health 2016;106(11):2042-8.
- 22. De Vaus DA. Surveys in social research. 6th ed. London: Routledge; 2014.
- 23. Townsend P. Social surveys of old age in Great Britain, 1945–58. Bull World Health Organ 1959;21:583–91.
- 24. Bowling A. Ageing well: quality of life in old age. Maidenhead, UK: Open University Press; 2005.
- 25. Fricker Jr RD. "Sampling methods for online surveys," the SAGE handbook of online research methods. 2016. p. 195–217.